



## Clinical Practice Guideline Pain and Bleeding in Early Pregnancy

### Target Audience

Registered Nurses/Midwives

Student Midwife/Nurse under supervision where appropriate and patient permits

Medical staff in Women's Health Unit

Medical staff in ED and Rapid Assessment Discharge (RAD) team members.

### Purpose

This CPG aims to clarify the initial and further care given to women who are experiencing per-vaginal (PV) bleeding and abdominal pain in early pregnancy:

- a. **Prior and up to 15+6 weeks gestation who attend the Emergency Department.**
- b. **After 16+0 weeks' gestation onwards, who attend the Emergency Department.**

The purpose is:

- To guide the initial assessment, investigation, and management of the woman in a timely manner.
- To guide the referral process to Women's Health Unit for follow up in women with gestation  $\geq 16$  weeks
- To guide the referral process to EPPAS for follow up in women with gestation  $\leq 15+6$  weeks.
- To provide supportive and consistent counselling to women with pain and bleeding in early pregnancy.
- To address the physical and psychosocial needs of the woman and her family.

### Definitions used in this Guideline.

- **Early pregnancy:** pregnancy up to 15+6 weeks of pregnancy *for the purpose of this guideline*. Some guidelines usually define early pregnancy as up to 12+0 weeks gestation.
- **Last menstrual period (LMP):** the first day of the last menstrual period
- **Gravida:** total number of confirmed pregnancies regardless of fetal outcome.
- **Parity:** the number of births a woman had after 20 weeks gestation regardless of fetal outcome.
- **Expectant management:** a management approach in which treatment (pharmacological or surgical) is not administered, with the aim of seeing whether the condition will resolve naturally.
- **Recurrent miscarriage:** Three or more consecutive miscarriages.
- **Threatened miscarriage:** Early viable intra-uterine pregnancy on ultrasound with clinical evidence of mild to moderate vaginal bleeding and closed cervical os. Abdominal pain may be present or not.
- **Incomplete miscarriage:** Sonographic presence of retained pregnancy tissue in utero measuring  $\geq 20$ mm with clinical evidence of a bulky uterus, vaginal bleeding

PROMPT doc no: 124294	Version: 6.0	
First created: 10/07/2015	Page 1 of 10	Last reviewed: 24/09/2020
Version changed: 24/09/2020	UNCONTROLLED WHEN DOWNLOADED	Next review: 24/09/2023



## Clinical Practice Guideline Pain and Bleeding in Early Pregnancy

and open cervix. Abdominal pain may be present or not. Sometimes products of conception may be seen in the cervix.

- **Missed miscarriage:** refers to a woman who has an ultrasound-defined non-viable pregnancy without pain or bleeding i.e. failure to identify an embryo in a gestational sac with MSD of  $\geq 25\text{mm}$  and or failure to identify cardiac activity in an embryo with CRL  $\geq 7\text{mm}$
- **Inevitable miscarriage:** applies to a woman who is pregnant and bleeding, with an open cervix. Sometimes products of conception can be visualized at the cervix.
- **Spontaneous miscarriage:** is defined a spontaneous and complete pregnancy loss prior to 20+0 weeks gestation.
- **Early pregnancy and Perinatal Assessment Service (EPPAS):** Outpatient service staffed by midwives, HMOs, GP Obstetricians with direct access to a consultant gynaecologist daily within normal working hours on Mondays to Fridays 0900 to 1630 hours. It is located in Outpatient Building 1. It is for women with non-acute concerns about early pregnancy pain and bleeding who are haemo-dynamically stable.
- **Heavy vaginal bleeding:**  $> 2$  soaked pads per hour.
- **Cervical shock:** a woman has vaginal bleeding and is hypotensive. Products of conception in the cervix can cause cervical dilation, vagal stimulation and vasovagal syncope. Typically, there is hypotension out of proportion to observed blood loss. Removal of the POC from the cervix allows recovery of the woman.
- **Pregnancy of unknown location (PUL):** a woman who has a positive  $\beta$ -hCG result but an ultrasound that cannot locate the site of pregnancy or identify RPOC. Ectopic pregnancy is not excluded. It may be an early pregnancy  $< 6$  weeks gestation, early failing pregnancy or complete spontaneous miscarriage.
- **Ectopic pregnancy:** a pregnancy located outside of the uterine cavity.
- **Heterotopic pregnancy:** two pregnancies where one sac is located within the uterine cavity and the other outside the uterine cavity
- **Products of conception (POC):** pregnancy related (fetal or placental) tissue including fetus, gestational sac, placenta, and decidua. This term is commonly used amongst health professionals, but women may prefer the term 'retained pregnancy tissue'.
- **Ovarian hyper stimulation syndrome (OHSS):** This can occur when ovulation stimulation medications are used in infertility management. Women can present with abdominal pain due to fluid accumulation in the abdomen and ovarian complications. These women are at risk of fluid overload (e.g. pulmonary edema), electrolyte disturbances, VTE and ovarian complications.

PROMPT doc no: 124294	Version: 6.0	
First created: 10/07/2015	Page 2 of 10	Last reviewed: 24/09/2020
Version changed: 24/09/2020	UNCONTROLLED WHEN DOWNLOADED	Next review: 24/09/2023



## Clinical Practice Guideline Pain and Bleeding in Early Pregnancy

### Background

Abdominal pain and bleeding are common in early pregnancy. Approximately 20% of pregnancies miscarry, and early pregnancy complications in the first trimester of pregnancy such as miscarriage and ectopic pregnancy can cause considerable distress.

Treatment and care should take into account women's needs and preferences. Women should have the opportunity to make informed decisions about their care and treatment, in partnership with their healthcare professional.

The major concern is the possibility of ectopic pregnancy, which can be life-threatening with an incidence of 1 in 200 – 500 pregnancies. It must be actively excluded in all women presenting with undifferentiated abdominal pain and/or per-vaginal bleeding with a positive pregnancy test.

Ectopic pregnancy should also be considered after a recent miscarriage or termination of pregnancy where no prior intrauterine gestational sac was identified on ultrasound, as heterotopic pregnancies can occur.

Common causes of abdominal pain may include:

- Ectopic pregnancy
- Miscarriage
- Ovarian pathology (e.g. cyst rupture, ovarian torsion, bleeding into a corpus luteum cyst)
- Genital tract infection
- Ovarian Hyper stimulation Syndrome OHSS.
- Non-gynaecological causes such as urinary tract infection, appendicitis and other gastro-intestinal tract disorders.

Common causes of bleeding may include:

- Miscarriage
- Retained Products of Conception RPOC
- Subchorionic haemorrhage
- Ectopic pregnancy
- Genital tract infection including endometritis
- Septicaemia which can occur secondary to infected RPOC and requires prompt management.
- Gestational trophoblastic disease
- Lower genital tract pathology such as cervical polyps, cervical erosions

### Support and Information Giving

- It is important to treat all women having early pregnancy complications with dignity and respect. Clinicians should be aware that women and their partners would react to complications or the loss of a pregnancy in different ways.
- Provide all women with information and support in a sensitive manner, taking into account their individual circumstances and emotional response.

PROMPT doc no: 124294	Version: 6.0	
First created: 10/07/2015	Page 3 of 10	Last reviewed: 24/09/2020
Version changed: 24/09/2020	UNCONTROLLED WHEN DOWNLOADED	Next review: 24/09/2023



## Clinical Practice Guideline Pain and Bleeding in Early Pregnancy

- Where possible, women should be cared for in appropriate rooms with consideration for privacy.
- Further care and support should be offered to these women via the social worker, SANDS <https://www.sands.org.au/>, or The Pink Elephants Support Network <https://miscarriagesupport.org.au/>.
- Throughout a woman's care, give her and (with agreement) her partner specific evidence-based information in a variety of formats on
  - When and how to seek help if existing symptoms worsen or new symptoms develop, including a 24-hour contact telephone number.
  - Where to access support and counselling services, including leaflets (EPPAS brochure), web addresses and helpline numbers for support organisations.

### Guideline

#### All women in early pregnancy

- a. prior and up to 15+6 weeks gestation and
- b. equal to or after 16+0 weeks gestation (unscheduled presentations)

who present with vaginal bleeding and abdominal pain to the Emergency Department must first be assessed and triaged by a triage competent nurse as appropriate, according to the Australian triage Scale ATS and their main presenting complaint.

#### Please note:

**Women who are  $\geq 16+0/40$  weeks gestation with a primary obstetric complaint e.g. labour or ruptured membranes will be asked to contact Women's Health triage midwife on 9784 7959 and be seen directly on Women's health unit.**

**Women who are  $\geq 16+0/40$  weeks gestation presenting with non-pregnancy related primary medical problems e.g. chest pain, asthma, shortness of breath should be seen and cleared initially by Frankston ED following which they should be reviewed/admitted in the Women's health unit.**

If the woman is an ATS Category 1 or 2, they must proceed immediately to the main department, to a resuscitation cubicle to receive prompt care.

If the woman is in ATS Categories 3, 4 or 5, then the triage nurse should:

- Take a set of vital signs to determine if patient is stable or unstable
- Assess pain and bleeding to assess severity of symptoms
- Request review by EDMO in fast track

PROMPT doc no: 124294	Version: 6.0	
First created: 10/07/2015	Page 4 of 10	Last reviewed: 24/09/2020
Version changed: 24/09/2020	UNCONTROLLED WHEN DOWNLOADED	Next review: 24/09/2023


# Clinical Practice Guideline Pain and Bleeding in Early Pregnancy

## Flowchart A: Assessment Algorithm: Pain and Bleeding in ED ≤ 15+6/40 weeks



## Clinical Practice Guideline Pain and Bleeding in Early Pregnancy

### Flowchart B: Assessment Algorithm: Pain and Bleeding in ED ≥ 16+0/40 weeks



PROMPT doc no: 124294	Version: 6.0
First created: 10/07/2015	Page 6 of 10
Version changed: 24/09/2020	UNCONTROLLED WHEN DOWNLOADED

Last reviewed: 24/09/2020

Next review: 24/09/2023



## Clinical Practice Guideline Pain and Bleeding in Early Pregnancy

### Anti-D administration.

- Anti D immunoglobulin (Ig) is used to inactivate fetal Rh-positive cells which may have crossed the placental barrier to enter the circulation of a Rh-negative mother.
- It is a blood product therefore written consent is required.
- Administration should be clearly documented in the maternal records and discharge summary (time and dose to be recorded in the medical notes).

### ADMINISTRATION

- A current group & hold should be initially taken or available to confirm Rh status.
- Blood bank should be contacted so Anti D Ig is issued.
- Following a sensitising event (confirmed miscarriage, ectopic pregnancy, invasive procedure), Anti D Ig should be administered as soon as possible and always within 72 hours to all Rh-negative women.
- Exceptionally, if this deadline is not met, some protection may still be offered if Anti D Ig is given up to 10 days after the sensitising event.
- Routine testing for feto-maternal haemorrhage is **not** required.

### DOSAGE

- 250 IU of Anti D Ig should be given where required at gestations  $\leq$  12 weeks.
- 625 IU of Anti D Ig should be given where required at gestations  $>$  12 weeks.
- 625 IU of Anti D Ig should be given for multiple pregnancy regardless of gestation.

### INDICATIONS FOR USE

- All miscarriages over 12 week's gestation including threatened miscarriage.
- All miscarriages where the uterus has been evacuated – medically or surgically.
- Threatened miscarriage particularly with gestation approaching 12 weeks if the bleeding has been heavy, repeated or associated with pain.
- Ectopic pregnancy whether medical or surgically managed.

### EXCLUSIONS TO USE

- Threatened miscarriage under 12 weeks.
- Complete miscarriage under 12 weeks where there has been no formal attempt to evacuate the uterus, as the risk of feto-maternal haemorrhage is negligible.

PROMPT doc no: 124294	Version: 6.0	
First created: 10/07/2015	Page 7 of 10	Last reviewed: 24/09/2020
Version changed: 24/09/2020	UNCONTROLLED WHEN DOWNLOADED	Next review: 24/09/2023



## Clinical Practice Guideline Pain and Bleeding in Early Pregnancy

### Management of POC (pregnancy tissue).

- If POC are spontaneously passed regardless of gestation, please contact the gynaecology registrar, as further management is required.
- POC can be gently assessed for completeness, the fetus and placenta may be identifiable.
- Please keep them aside fresh (not in formalin) and as undisturbed as possible. They may need to be reviewed by the Gynaecology Registrar and be sent for histology or karyotyping, particularly if history of recurrent miscarriage.
- POC should be kept in the same area as the mother, but can be obscured from view or left in an adjacent room if distressing.
- Some mothers may choose to keep the POC for burial or grieving – this can be facilitated and staff should consider consulting RAD for further information about such services.

### Comments about $\beta$ -hCG.

- $\beta$ -hCG levels must be considered in the context of the clinical picture. Interpretation can be difficult and values often have wide ranges.
- Gestational age assessment should be done based on ultrasound and not  $\beta$ -hCG levels.
- It is important that counselling given to women with early pregnancy complications is **not** provided on the basis of  $\beta$ -hCG levels alone.

After approximately 7-week's gestation, short-term changes in  $\beta$ -hCG levels are unlikely to be solely helpful, in interpreting the clinical picture (with the exception of catastrophic falls).

$\beta$ -hCG levels peak in the late first trimester around 11 to 12 weeks, after which they plateau and fall in normal pregnancy. hCG levels peak in the late first trimester around 11 to 12 weeks, after which they plateau and fall in normal pregnancy. fall in normal pregnancy hCG levels

PROMPT doc no: 124294	Version: 6.0	
First created: 10/07/2015	Page 8 of 10	Last reviewed: 24/09/2020
Version changed: 24/09/2020	UNCONTROLLED WHEN DOWNLOADED	Next review: 24/09/2023



## Clinical Practice Guideline Pain and Bleeding in Early Pregnancy

peak in the late first trimester around 11 to 12 weeks, after which they plateau and fall in normal pregnancy.

**It is therefore not recommended to routinely perform serum  $\beta$ -hCG after 12 weeks gestation.**

It is best that interpretation of  $\beta$ -hCG trends is performed by the EPPAS team or Gynaecology Registrar.

As a general guide:

- An early pregnancy is identifiable on transvaginal pelvic ultrasound if the  $\beta$ -hCG level is >1500 IU/L (discriminatory  $\beta$ -hCG zone).
- Fetal cardiac activity is usually seen after 6 weeks gestation or when the  $\beta$ -hCG level is >10,000 IU/L
- 66% of normal viable intrauterine pregnancies will demonstrate doubling of  $\beta$ -hCG levels double every 48 hours in the first 8 weeks of pregnancy.
- 15% of viable intrauterine pregnancies will not demonstrate a normal  $\beta$ -hCG doubling time in the first 7 weeks of pregnancy.
- 13% of ectopic pregnancies will also demonstrate a normal  $\beta$ -hCG doubling time.
- A sustained fall in  $\beta$ -hCG may be suggestive of miscarriage. It does not exclude ectopic pregnancy.
- A plateau or slow rise/fall in  $\beta$ -hCG may be suggestive of either a miscarriage or ectopic pregnancy.

### Key Aligned Documents

- Management of Ectopic Pregnancy
- Anti – Rh (D) Immunoglobulin antenatal and postnatal administration
- Blood and Body Substance Exposure Prevention and Management of Exposure
- Hand Hygiene & Aseptic Technique
- Initial management of critically ill obstetric patient in ED/Labour ward
- Postpartum haemorrhage

### References

1. NICE guideline. Ectopic pregnancy and miscarriage: diagnosis and initial management. Published 17 April 2019. [www.nice.org.uk/guidance/ng126](http://www.nice.org.uk/guidance/ng126)
2. RCOG guideline. Diagnosis and management of ectopic pregnancy. Green-top guideline No. 21. Published 04 April 2016. [www.rcog.org.uk/en/guidelines-research-services/guidelines/gtg21/](http://www.rcog.org.uk/en/guidelines-research-services/guidelines/gtg21/)

PROMPT doc no: 124294	Version: 6.0	
First created: 10/07/2015	Page 9 of 10	Last reviewed: 24/09/2020
Version changed: 24/09/2020	UNCONTROLLED WHEN DOWNLOADED	Next review: 24/09/2023

## Clinical Practice Guideline Pain and Bleeding in Early Pregnancy

3. RCOG Guideline. The Investigation and Treatment of Couples with Recurrent First-trimester and Second-trimester Miscarriage. Green-top guideline No. 17. Published April 2011. Updated Feb 2017. <https://www.rcog.org.uk/en/guidelines-research-services/guidelines/gtg17/>

Document management	Position
Executive Sponsor:	Executive Director Medical Services Frankston Hospital
Document Author	Clinical Director Women's Health
Document Owner:	Clinical Director Women's Health
Date approved	August 2020
Approved by:	Emergency Department Working Group
Date created/revised in archived system:	02/2011, 05/2017, 05/2019, 09/2020

PROMPT doc no: 124294	Version: 6.0	
First created: 10/07/2015	Page 10 of 10	Last reviewed: 24/09/2020
Version changed: 24/09/2020	UNCONTROLLED WHEN DOWNLOADED	Next review: 24/09/2023